

Trends

in Psychiatry and Psychotherapy

JOURNAL ARTICLE PRE-PROOF **(as accepted)**

Review Article

Blended Psychological Interventions for Emotional Disorders in Youth: Acceptability and Intention to Use in a Sample of Portuguese Psychologists

Bárbara Gomes-Pereira, Ana Góis, Ana Maria Pereira, Brígida Caiado, Helena Moreira, Ana Isabel Pereira

<http://doi.org/10.47626/2237-6089-2023-0770>

Original submitted Date: 28-Nov-2023

Accepted Date: 13-Jun-2024

This is a preliminary, unedited version of a manuscript that has been accepted for publication in Trends in Psychiatry and Psychotherapy. As a service to our readers, we are providing this early version of the manuscript. The manuscript will still undergo copyediting, typesetting, and review of the resulting proof before it is published in final form on the SciELO database (www.scielo.br/trends). The final version may present slight differences in relation to the present version.

Blended Psychological Interventions for Emotional Disorders in Youth: Acceptability and Intention to Use in a Sample of Portuguese Psychologists

Short title: Blended Psychotherapy for Youth Emotional Disorders

Bárbara Gomes-Pereira¹, Ana Góis¹, Ana Maria Pereira², Brígida Caiado¹,
Helena Moreira¹, Ana Isabel Pereira²

¹University of Coimbra, Centre for Research in Neuropsychology and Cognitive Behavioral Intervention, Faculty of Psychology and Educational Sciences, Coimbra, Portugal.

²University of Lisbon, Faculty of Psychology, Lisboa, Portugal.

Corresponding Author:

Bárbara Gomes-Pereira

Colégio Novo, 3000-115 Coimbra

Coimbra

3000-115

barbarapereirapsi@outlook.com

Abstract

Introduction: The rise of mental health problems in youth highlights the need for accessible and cost-effective psychological interventions. Blended interventions, which combine face-to-face and online sessions, can be an adequate response to the increase in demands for youth mental health services. Although this can be a promising approach, effective dissemination depends on the professionals' acceptance.

Objectives: This study aimed to explore the acceptability of and intention to use blended interventions by psychologists working with children with emotional disorders and to examine their predictors, including previous knowledge,

expectancies (i.e., performance expectancy, effort expectancy, social influence, and facilitating conditions), and attitudes toward evidence-based practices.

Methods: The sample consisted of 76 Portuguese psychologists (Mage = 37.26 years, SD = 10.47; 92.1% female) working in youth mental health services. The participants completed an online protocol to evaluate the different dimensions included in the study.

Results: The results showed that most participants demonstrated moderate to high acceptability of blended psychological interventions for emotional disorders in youth and intend to use them in the future. Regression analysis showed that performance expectancy and positive attitudes toward evidence-based practices were significant predictors of acceptance of blended interventions and that social influence was a significant predictor of both acceptance of and intention to use blended interventions.

Conclusion: These results emphasize the importance of sharing the findings of blended interventions, changing professionals' attitudes toward evidencebased practices, and collaborating more closely with organizations and institutions to advance standards that encourage the adoption of this intervention format.

Keywords: Acceptance; Attitudes and expectancies; Blended interventions; Intention to use; Psychologists; Youth emotional disorders.

Introduction

Mental health problems in children and adolescents represent a growing public health concern because of their long-lasting negative effects.¹ Specifically, emotional disorders (i.e., a term that groups together anxiety disorders, anxiety-related disorders and depressive disorders)^{2,3} have a negative effect on children's development^{4,5} and quality of life⁶ and impact different domains of child life, contributing to poor academic performance or social functioning.⁷ In fact, the meta-analysis by Polanczyk et al.,⁸ conducted with data from 27 countries reported a worldwide prevalence of 6.5% for any anxiety disorder and 2.6% for any depressive disorder in youth, and recent studies suggest that the COVID-19 pandemic has increased the prevalence of these problems in youth.⁹

Online and Blended Psychological Interventions for Youth

Despite widespread recognition of the importance of mental health promotion and prevention in children and adolescents, an enormous disparity between the available resources and actual access to mental healthcare remains.^{10,11} Among the obstacles to obtaining proper mental health care are low socioeconomic resources, stigma, and accessibility problems,¹² such as geographic distance to mental health services, lack of time (of patients and professionals), long waiting times, and service costs.^{13,14} One may argue that relying only on conventional intervention delivery methods is insufficient, and online interventions may help overcome these obstacles.

In online psychological interventions, participants are not required to travel to sessions. They may also feel that their privacy is protected with this type of intervention, allaying any concerns about the stigma associated with face-to-face therapy.¹⁵ Previous research in adult populations has shown that online cognitive behavioral therapy (CBT), especially with therapist guidance, is efficacious when used for the treatment of emotional disorders^{16,17} (e.g., depression). Some self-guided interventions for children with anxiety symptoms have also been developed, such as Lumi Nova (BfB labs)¹⁸ and BRAVE-online.¹⁹ Research has shown that these interventions potentially reduce symptoms,^{20,21} and have been accepted by children, parents and clinicians.²² However, several limitations emerge in online self-guided interventions, such as the absence of human contact, the unreliability and failure of technological equipment, limited internet access, or the need for more participant autonomy since there are no face-to-face sessions to clarify questions that may arise.²³

A blended format that combines online sessions and face-to-face sessions can overcome some of the limitations of self-guided online interventions, enabling a more tailored intervention to a child's needs and developing a relationship between a patient and therapist through the face-to-face sessions²⁴. The blended format simultaneously promotes cost-effectiveness since it maintains some of the characteristics of the online format, allowing for greater accessibility to treatment.²⁵ Therefore, blended therapy is suggested as a promising innovation for the psychotherapeutic setting.²⁶

Predictors of Acceptability and Intention to Use Blended Psychological Interventions for Children with Emotional Disorders

One of the most critical factors in determining whether patients use blended programs is the therapists' acceptability of this type of delivery format, which may be described as "the extent to which people delivering or receiving a healthcare intervention consider it to be appropriate, based on anticipated or experienced cognitive and emotional responses to the intervention".²⁷ Some studies aiming to examine attitudes toward blended therapy show that blended treatment is generally accepted, although psychotherapists do not prefer web-based or blended therapy over face-to-face therapy.^{28,29} To the best of our knowledge, in Portugal, only one published study has addressed psychologists' attitudes toward online psychological interventions for adults. Mendes-Santos et al.,³⁰ found that most Portuguese psychologists had a slightly negative/neutral view toward such treatments and were unfamiliar with them, had no specific training, and had no prior experience utilizing online therapies. **These results in Portugal starkly contrast with countries such as Australia,³¹ the United Kingdom, and Sweden,³² where the use of internet interventions is widely disseminated. As conceptualized by Topooco et al,³² Portugal may be included in the 'learners' category in this domain, given the very limited current experience and practice of e-mental health in the country.** In the latter study of Mendes-Santos et al.,³⁰ the authors also found greater acceptability toward blended treatment interventions compared to self-guided online interventions.

Attitudes toward manualized evidence-based treatment (EBT) for adults might also be an important determining factor of the acceptability of online and blended interventions by professionals,³³⁻³⁵ which are usually structured and manualized. In general, there are still negative attitudes toward manualized EBTs, with several professionals perceiving them as less relevant to their clinical work than other factors³⁶ (e.g., clinical experience) and not valuing, or only minimally valuing, the role of research on their clinical practice.³⁷ According to earlier research, professionals' negative attitudes toward web-based solutions were generally recognized as barriers to their effective uptake and recommendation.^{33,38} However, despite this evidence, there has been a growing movement toward acceptance of EBPs. For example, the study by Lilienfeld et

al.,³⁹ showed that many or most mental health professionals had a reasonably positive view of evidence-based practices (EBPs) and their usefulness in clinical practice. There is also evidence that therapists have an interest in and positive attitudes toward the implementation of blended therapy³⁵ in particular, which is considered to be a facilitator to the uptake of these interventions.

Other factors, such as therapists' knowledge of online and blended treatments and their prior usage,³⁰ were proposed as potential predictors of acceptability and whether therapists will use online therapy. Specifically, Mendes-Santos et al.,³⁰ pointed to the lack of knowledge and training as one of the main obstacles to overcome in order to guarantee the successful implementation of online interventions.

To better understand the predictors of users' intentions to use information technology (IT), such as an online psychological intervention, and their subsequent usage behavior, the Unified Theory of Acceptance and Use of Technology (UTAUT) was developed.⁴⁰ The UTAUT is based on the theory of reasoned action (TRA), the theory of planned behavior (TPB), and the social cognitive theory (SCT).⁴¹ According to this model, four cognitive dimensions may play a significant role as direct determinants of the intention to use an online intervention: performance expectancy (i.e., how much a person thinks the intervention will work and be beneficial); effort expectancy (i.e., the degree to which a person believes that adopting the intervention will be easy); social influence (i.e., how much a person believes that others believe they should use the intervention); and facilitating conditions (i.e., the degree to which an individual believes that an organizational and technical infrastructure exists to support the use of the system).

However, to the best of our knowledge, all the studies conducted on the acceptance and intention to use blended interventions by psychologists have only been carried out as a whole and not specifically for use with children. In other words, there is a need to understand whether professionals' levels of acceptance and intention to use, as well as the predictors of these variables, change when the target population of the blended intervention becomes children with emotional disorders.

The Present Study

In this study, we intended to explore the predictors of acceptability and intentions of Portuguese psychologists to use blended interventions for children with emotional disorders. Specifically, we aimed to (1) describe psychologists' prior knowledge and experience with blended interventions, as well as their level of acceptability of and intention to use these interventions, and (2) assess the role that psychologists' expectations and attitudes toward manualized evidence-based practices may have on their acceptability of and intention to use blended interventions for children with emotional problems.

Method

Participants

The study sample comprised 76 Portuguese psychologists ($M_{age} = 37.26$ years, $SD = 10.47$; 92.1% female). Most participants reported having a bachelor's or master's degree (88.2%), having a clinical and health psychology specialization (73.7%), and adopting a cognitive-behavioral approach (60.5%). The detailed characteristics are presented in Table 1.

Table 1 - Sociodemographic, academic, and professional background variables

	<i>n</i>	%
Gender		
Female	70	92.1
Male	6	7.9
Professional work location		
Urban	63	82.9
Rural	13	17.1
Academic training		
Bachelor (5 years) or Master	67	88.2
PhD	9	11.8
Specialization		

	<i>n</i>	%
Clinical and Health Psychology	56	73.7
Education Psychology	17	22.4
Organizational Psychology	1	1.3
Junior Psychologists	5	6.6
No Specialization	7	9.2
Theoretical Approach		
Cognitive-behavioral	46	60.5
Psychodynamic	10	13.2
Systemic	7	9.2
Integrative	9	11.8
Humanist	3	3.9
Other	1	1.3
Years of professional experience		
0 – 3 years	23	30.3
4 - 15 years	26	34.2
16 - 37 years	27	35.5
Practice Context		
Central Hospital	6	7.9
Private Hospital	1	1.3
Primary Care Center	5	6.6
Private Practice	29	38.2
School	19	25.0
Other	16	21.1

Procedures

Ethical approval for the current study was obtained from the Ethics and Deontology Committee of the Faculty of Psychology of the University of Lisbon and University of Coimbra (CEDI/23/06/2021). **The inclusion criteria were being a psychologist working in the field of child mental health (including psychologists**

working with adolescents or parents). All participating professionals completed the questionnaires via a data collection website (LimeSurvey®) between June 2021 and March 2022. The study and the survey link were shared on social media, on the website of the Portuguese Psychologists' Association and via email. Before starting the study, participants were informed about the definition of blended interventions (i.e., which combine online and face-to-face sessions), informed about the main objectives of the study and the anonymity of their responses. After reading the information about the study, participants had the option of going ahead and giving their consent to participate. A total of 220 psychologists started answering the online questionnaire, but only 76 completed it.

Measures

Sociodemographic questionnaire

The first part of the questionnaire, which was developed based on previous sociodemographic questionnaires used by our research team, concentrated on gathering essential sociodemographic information. Participants were asked about age, sex, nationality, academic background, specialization in psychology, advanced specialization in psychology, main theoretical orientation of intervention, professional activities performed, context and location of professional activities, district (region) of the workplace and number of years of professional experience.

Previous experience with online and blended interventions

Psychologists were asked several questions regarding their previous experience with online and blended interventions, including questions such as "In your professional practice, have you ever implemented any online, or blended psychological intervention program for children or adolescents?"; "In your professional practice, do you usually recommend or have you ever recommended the use/consultation of online resources to children, adolescents, or parents as a complement to the psychotherapeutic process?"; and "Which online resources do you usually recommend or have you ever recommended?".

Degree of knowledge of online and blended interventions

A short scale to evaluate the psychologists' level of knowledge about several aspects of online and blended interventions was developed for the current study. The scale was composed of six items, rated on a Likert scale ranging from 1 ("nonexistent") to 5 ("high"). The first three questions assessed knowledge about the content, mode of functioning, and existing research regarding self-guided online interventions (regardless of age group), and the remaining three questions asked the same questions regarding blended interventions. A single factor emerged from the six items as a result of a principal component analysis. Cronbach's alpha for the total score was .96).

Attitudes Toward Manualized Evidence-Based Practice

The Portuguese version of the Evidence-Based Practice Attitude Scale (EBPAS)⁴⁰ was used to assess psychologists' attitudes toward manualized evidence-based treatments. Therefore, the following explanation was provided in the instructions of the questionnaire: "Manualized therapy refers to any intervention that has specific guidelines and/or components that are outlined in a manual and/or that are to be followed in a structured/predetermined way". The questionnaire includes 15 items, which are rated on a five-point Likert scale (0 = "strongly disagree" to 4 = "strongly agree"), with higher scores indicating more favorable attitudes. Cronbach's alpha for the total scale was .73.

Expectancies towards Blended Psychological Interventions for Children with Emotional Disorders

A questionnaire, based on the Unified Theory of Acceptance and Use of Technology (UTAUT)⁴⁰ model, was specifically developed for the current study. Principal component analysis was performed to examine the factor structure of the questionnaire. Four factors emerged from the analysis: Performance Expectancy (7 items assessing the degree to which an individual believes that using the system will help him or her to attain gains in job performance/perceived usefulness; e.g., "A blended psychological intervention would increase the effectiveness of my clinical work"); Social Influence (4 items assessing the degree to which an individual perceives that significant others believe he or she should

use the system/subjective norms; e.g. “My superiors would support my decision to implement a blended psychological intervention”); Facilitating Conditions (7 items assessing the degree to which an individual believes that an organizational and technical infrastructure exists to support the use of the system/perceived behavioral control; e.g. “The institution where I work would provide the necessary means for me to implement the intervention”); and Effort Expectancy (2 items assessing degree of ease associated with the use of the system; e.g., “Implementing a blended intervention would require too much of my time and energy”). The items are rated on a Likert scale ranging from 1 (“strongly disagree”) to 5 (“strongly agree”). Cronbach's alpha reliabilities were .82 for Performance Expectancy; .85 for Social Influence; .70 for Facilitating Conditions; and .65 for Effort Expectancy.

Perceived Acceptability of Blended Psychological Interventions for Children with Emotional Disorders

A 3-item questionnaire was developed for this study to assess the psychologists' perceived usefulness of blended psychological interventions for children with mild and severe anxiety and/or mood problems (e.g., “To what extent do you think blended psychological interventions can help children aged 6 to 13 years with moderate anxiety or mood problems?”). Items were rated on a five-point Likert scale ranging from 1 (*not at all*) to 5 (*very*). Cronbach's alpha was found to be good (.80).

Intention to Use Blended Psychological Interventions for Children with Emotional Disorders

The intention of psychologists to use or recommend blended psychological interventions for children with emotional disorders was evaluated by two items: “If an empirically validated blended psychological intervention was available for children aged 6 to 13 years with anxiety or mood disorders, would you consider using it in your professional practice?” and “If an empirically validated blended psychological intervention was available for children aged 6 to 13 years with anxiety or mood disorders, would you consider recommending it to a colleague?”.

These two items were answered on three-point Likert scale (0 = "no"; 1 = "maybe"; 2 = "no") and has a Cronbach's alpha of .87.

Data Analyzes

For the data analysis, we used the Statistical Package for the Social Sciences (SPSS) statistical analysis software, version 28, for Windows. Descriptive analyzes were performed to characterize the sample regarding sociodemographic dimensions and to evaluate the degree of knowledge, experience levels, and preferences of the professionals concerning online and blended psychological interventions. The reliability of the various questionnaires used was examined through Cronbach's alpha, where values above .70 are recommended.⁴³ Correlations between all the variables under study were also checked. Two stepwise regression analyses were performed to examine the predictors of 1) psychologists' acceptability toward blended interventions and 2) psychologists' intention to use these types of interventions. Years of professional experience were input into the first step, degree of knowledge was added to the second step, EBPAS was included in the third step, and lastly, the four UTAUT model elements were added in the final step of the model.

Results

Experience and Knowledge about Online or Blended Psychological Interventions

The mean values and standard deviations of knowledge regarding self-guided online and blended psychological interventions are presented in Table 2.

Table 2 - Mean values of knowledge about online or blended psychological interventions

	<i>M</i>	<i>SD</i>
Self-guided Online Psychological Interventions		
Content	2.41	1.21
How it works	2.49	1.24
Research	2.20	1.2
Blended Psychological Interventions		
Content	2.51	1.21

How it works	2.64	1.28
Research	2.39	1.28

Note. Items of the presented scale assume Likert values between 1 ("nonexistent") and 5 ("high").

Most professionals did not know of online or blended psychological intervention programs for children or adolescents (93.4%). None of the 76 participants had ever implemented online or blended psychological intervention programs for children or adolescents.

In their professional practice, 34 (44.7%) out of the 76 participants had already recommended the use/consultation of online resources (e.g., websites, forums, social networks) to children, adolescents, or parents to complement the therapeutic process. The most recommended resources were websites related to mental health or another specific theme ($n = 29$, 38.2%). In addition, most professionals (88.2%) had used videoconferencing programs (e.g., Skype, Zoom, Teams) to conduct consultations.

Preliminary Results

The majority of professionals said they would consider using a blended intervention in their professional practice if it were available and would recommend it to a colleague (57.9% and 64.5%, respectively). All correlations between the predictors of the intention to use and acceptability of blended interventions and the levels of acceptance by professionals are presented in Table 3.

Table 3 - Scale correlations and means

	1	2	3	4	5	6	7	8	9
1. Years of Professional Experience	-								
2. Degree of Knowledge Total	-.11	-							
3. EBPAS Total	-.23*	.08	-						
4. Performance Expectancy	-.16	.10	.57**	-					

5. Social Influence	-.04	.28*	.45**	.45**	-				
6. Facilitating Conditions	.04	.17	.36**	.40**	.50**	-			
7. Effort Expectancy	-.04	.09	-.14	-.06	-.21	-.05	-		
8. Acceptance	-.20	.16	.65**	.62**	.58**	.39**	-.11	-	
9. Intention to use	.01	.21	.38**	.38**	.49**	.40**	-.10	.56**	-
Scale M (SD)	11.5 (9.98)	14.64 (6.75)	2.92 (.46)	3.62 (.68)	3.49 (.68)	3.68 (.67)	2.80 (.76)	3.55 (.70)	1.59 (.50)

Note. * $p < .05$ *** $p < .001$; EBPAS = Evidence-Based Practice Attitude Scale; M(SD) = Mean (Standard Deviation);

Predictors of Acceptance and Intention to Use Blended Psychological Interventions for Children with Emotional Disorders

Two stepwise regression analyses were performed to assess the prediction of variables related to the acceptability of and intention to use blended interventions. Regarding the perceived acceptability of blended interventions, the results showed significant effects of social influence and performance expectancy, meaning that higher levels of these dimensions were predictive of higher levels of acceptability. The results also revealed that attitude toward evidence-based practice was a significant predictor of the acceptability of blended interventions (see Table 4).

Table 4 - Summary Stepwise Regression Analyzes for the Acceptability of Blended Interventions

Variable	Step 1			Step 2			Step 3			Step 4		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Years of Professional Experience	-.01	.01	-.20	-.01	.01	-.18	-.00	.01	-.04	-.00	.01	-.06
Degree of Knowledge				.02	.01	.14	.01	.01	.11	.00	.01	.02
EBPAS total score							.98	.14	.64**	.54	.16	.35*
Performance Expectancy										.29	.10	.28*

Social Influence				.29	.11	.28*
Facilitating Conditions				.02	.10	.02
Effort Expectancy				.01	.08	.01
<i>F Change</i>	2.98	1.59	49.17***	5.74***		
<i>R² Change</i>	.04	.02	.38	.14		

Note. ** $p < .01$ *** $p < .001$; EBPAS = Evidence-Based Practice Attitude Scale; SE B = Standard Error of the Coefficient B.

With regard to the intention to use blended interventions, the only significant predictor was social influence (see Table 5).

Table 5 - Summary Regression Analyses for the Intention to Use Blended Interventions

Variable	Step 1			Step 2			Step 3			Step 4		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Years of Professional Experience	.00	.01	.01	.00	.01	.03	.01	.01	.12	.00	.01	.08
Degree of Knowledge Total				.02	.01	.22	.02	.01	.20	.01	.01	.10
EBPAS Total							.43	.12	.39***	.16	.14	.15
Performance Expectancy										.09	.10	.13
Social Influence										.20	.10	.26*
Facilitating Conditions										.11	.09	.15
Effort Expectancy										-.01	.07	-.02
<i>F Change</i>	.00			3.53			13.0***			3.02*		
<i>R² Change</i>	.00			.05			.15			.12		

Note. * $p < .05$ *** $p < .001$; EBPAS = Evidence-Based Practice Attitude Scale; SE B = Standard Error of the Coefficient B.

Discussion

The primary objective of this study was to investigate the level of acceptability of and intention to use blended psychological interventions for treating emotional disorders in children among Portuguese psychologists and identify the role of psychologists' expectations and attitudes toward manualized evidence-based practices on these two variables.

The results showed that most psychologists were not familiar with online or blended interventions for children and adolescents with emotional disorders, and none of them had experience implementing this type of intervention in their practice. These results align with previous studies (e.g., Mendes-Santos et al., 2020), and underline the need for more knowledge and experience of Portuguese psychologists in the use of online interventions.³⁰ As mentioned above, some authors even point out that Portugal may fall under the learner's group in this area due to the country's lack of online mental health knowledge and usage.³² However, in the current study, most of the participants had experience conducting online consultations. The fact that the sample was recruited during the COVID-19 pandemic may have influenced these results since the large-scale self-quarantine and shelter-in-place orders led many nonemergency medical services to adopt telehealth solutions to continue serving their patients.^{44,45}

In addition, it was possible to verify that the psychologists had moderate to high levels of acceptance and intentions to use a blended intervention for children between 6 and 13 years old with emotional disorders. These results are in line with other studies carried out in recent years^{46,47} and with studies carried out specifically in Portugal, in which, as previously mentioned, mental health professionals showed a high acceptance rate in relation to the use of blended therapies compared to fully online therapies.³⁰ It is important to note that none of these studies have been carried out specifically on the use of these interventions for children and that, to our knowledge, this study is innovative in this age group. Although several studies have pointed out that there is still a long way to go, the truth is that the opinion of therapists regarding the use of online interventions (mainly blended interventions) has been changing and becoming increasingly positive. Part of this change can be justified by the impact that the COVID-19 pandemic has had around the world. This hypothesis is supported by authors

such as Wind et al.,⁴⁸ who considered the outbreak of the pandemic to be a "turning point" for e-mental health since it increased the use of technologies for therapeutic purposes.

This study also explored potential predictors of psychologists' acceptability of blended interventions. The results showed that years of experience and the level of knowledge of blended interventions did not predict psychologists' acceptability, neither facilitating conditions nor effort expectancy. Nevertheless, positive attitudes toward manualized evidence-based practices, social influence (i.e., the degree to which a psychologist believes that significant others, such as work colleagues or superiors, consider that a blended strategy should be used to address children's emotional disorders), and performance expectancy (i.e., the degree to which a psychologist believes that using a blended intervention for children with an emotional disorder will be effective and useful in their clinical practice) were shown to be significant predictors of the acceptability of blended interventions. These results are in line with previous studies that point to performance expectancy and social influence as important predictors for increasing practitioners' acceptability of online interventions.^{40,49,50} For instance, Philippi et al.,⁵¹ reported that regarding practitioners' acceptability of using web-based and mobile interventions, the main predictor was indeed performance expectancy, and Venkatesh et al.,⁴⁰ suggested performance expectancy to be the most critical predictor of eHealth acceptance. Regarding social influence, these results are in line with those found in pediatric health care, which showed that the positive social influence of peers and parents had a significant positive effect on eHealth experiences^{49,50} and was identified as a facilitator to the acceptance of eHealth interventions.⁵²

The predictors of psychologists' intentions to use blended therapies were also explored in this study, and the only significant predictor was social influence. Considering other studies, this result does not agree with the research previously conducted, which points to social influence as a nonsignificant variable of behavioral intention and points to performance expectancy as the most significant predictor.^{53,54} The apparent inconsistency in the results of these studies may be attributed, in part, to variations within the analysed samples. Previous studies included both mental health counsellors and primary care psychologists,

professional groups whose realities can significantly differ from those encompassed by the current study. Consequently, the impact derived from peer encouragement might not manifest in the same manner, potentially altering the value of social influence. This incongruity can be elucidated by the disparities in the samples, which encompass differences not only in professional background but also in cultural aspects, thereby contributing to the observed variability in the results. In addition, it is also worth considering that the limited knowledge and experience demonstrated by Portuguese professionals in utilizing online and blended therapies^{30,44} might render their intentions to use such interventions more reliant on social factors. The relatively low familiarity and expertise within the realm of blended therapies could lead to a heightened dependency on social aspects when contemplating the adoption of this intervention approach.

The fact that social influence has been shown to be a significant variable in both acceptance and intention to use blended interventions suggests that it would be important for this variable to be targeted and for strategies to be developed that focus on it in order to increase it. The integration of specific training for blended interventions in professional or educational settings could be an effective way to increase psychologists' acceptability of these interventions.

Limitations

While this study contributes to the understanding of the acceptance of and intention to use blended interventions in a Portuguese context, it also has some limitations that should be considered. **The sample primarily consisted of cognitive-behavioral therapists (60.5%), thus caution is warranted when extrapolating the findings to all mental health professionals in Portugal, as certain areas may be underrepresented. Furthermore, the significant variation in years of professional experience, ranging from novice practitioners to those with no prior experience, may pose an additional limitation as it complicates direct comparisons.** In future studies, greater equality in the various theoretical perspectives of the professionals under study is crucial because this is a factor that may greatly influence their preferences. A second limitation stems from significant participant attrition during the online questionnaire, likely due to its extensive nature. This dropout pattern, observed from the outset, aligns with prior

research.⁵⁵ Future studies may benefit from adopting strategies proposed in existing literature, such as a two-phase approach: the first gathering consent, contact, and demographic data, and the second focusing on survey completion.⁵⁵ A third limitation relates to the fact that most of the measures used, although derived from the literature and demonstrating adequate reliability, were specifically adapted for this study. This adaptation may hinder the comparison of the results with existing literature. Additionally, particular attention should be given to the two measures that contain few items (e.g., 2 or 3 items), as they may lead to less reliable conclusions. One additional limitation is based on the fact that in contrast to the longitudinal nature of the original UTAUT study,⁴⁰ this research had a cross-sectional design. Given the scarcity of empirical research on UTAUT and technology adoption, it is imperative to conduct longitudinal studies to facilitate comparisons with Venkatesh's UTAUT⁴⁰ investigations and enhance the understanding of technology adoption and usage.

Conclusions

With the increasing number of mental health problems in children and adolescents and the difficulty in finding resources in the current healthcare system, it is essential to develop and disseminate new and cost-effective therapeutic solutions. A blended format combining online and face-to-face sessions can overcome some of these problems. This allows greater accessibility to treatment, is more adapted to the needs of the children, and encourages a closer therapeutic relationship. Therefore, evaluating the acceptability of and intention to use these interventions among mental health professionals, such as psychologists, as well as their predictors, is critical for better disseminating them among professionals and increasing the likelihood of usage.

Overall, this study showed that despite low knowledge and experience using online or blended interventions, psychologists' acceptability and attitudes toward these interventions are positive. In addition, the study showed that variables such as performance expectancy and social influence can predict higher levels of acceptance and intention to use. Based on these results and considering the positive levels of acceptance, it seems necessary to start training

mental health professionals on the use of these interventions and allow them to gain knowledge and experience in using them.

Author contributions: CRediT Taxonomy Bárbara Gomes-Pereira Conceptualization-Lead, Formal analysis-Lead, Investigation-Equal, Methodology-Lead, Software-Lead, Writing – original draft-Lead Ana Góis Conceptualization-Supporting, Data curation-Supporting, Resources-Supporting Ana Maria Pereira Conceptualization-Supporting, Data curation-Supporting, Resources-Supporting Brígida Caiado Conceptualization-Supporting, Data curation-Supporting, Resources-Supporting Helena Moreira Conceptualization-Supporting, Data curation-Supporting, Project administration-Lead, Supervision-Supporting, Writing – review & editing-Supporting Ana Isabel Pereira Conceptualization-Supporting, Data curation-Supporting, Formal analysis-Lead, Project administration-Lead, Resources-Lead, Supervision-Lead, Writing – review & editing-Lead

Total of words: 3884

Date of the last literature: 25.08.2023

This work was supported by the Portuguese Foundation for Science and Technology [grant number PTDC/PSI-GER/0689/2020]. This work was also supported by individual grants research by the Portuguese Foundation for Science and Technology given to the first three authors [2022.13843.BD; 2020.09015.BD 2020.06835.BD].

Handling Editor: Mrs. Joana Bücker

References

1. Kieling C, Baker-Henningham H, Belfer M, Conti G, Ertem I, Omigbodun O, et al. Child and adolescent mental health worldwide: evidence for action. *Lancet*. 2011;378:1515-25.
2. Barlow DH, Farchione TJ, Sauer-Zavala S, Murray Latin H, Ellard KK, Bullis JR, et al. Unified Protocol for Transdiagnostic Treatment of Emotional

Disorders Therapist Guide: Therapist Guide. New York: Oxford Univ Press; 2018.

3. Bullis JR, Boettcher H, Sauer-Zavala S, Farchione TJ, Barlow DH. What is an emotional disorder? A transdiagnostic mechanistic definition with implications for assessment, treatment, and prevention. *Clin Psychol Sci Pract.* 2019;26:20.
4. Cartwright-Hatton S, McNicol K, Doubleday E. Anxiety in a neglected population: Prevalence of anxiety disorders in pre-adolescent children. *Clin Psychol Rev.* 2006;26:817-33.
5. Kovess V, Carta MG, Pez O, Bitfoi A, Koç C, Goelitz D, et al. The School Children Mental Health in Europe (SCMHE) Project: Design and First Results. *Clin Pract Epidemiol Ment Health.* 2015;11(Suppl 1 M7):113-23.
6. Martinsen KD, Neumer SP, Holen S, Waaktaar T, Sund AM, Kendall PC. Self-reported quality of life and self-esteem in sad and anxious school children. *BMC Psychol.* 2016;4:45.
7. Fergusson DM, Woodward LJ. Mental health, educational, and social role outcomes of adolescents with depression. *Arch Gen Psychiatry.* 2002;59:225-31.
8. Polanczyk GV, Salum GA, Sugaya LS, Caye A, Rohde LA. Annual research review: A meta-analysis of the worldwide prevalence of mental disorders in children and adolescents. *J Child Psychol Psychiatry.* 2015;56:345-65.
9. Hawes M, Szenczy A, Klein D, Hajcak G, Nelson B. Increases in depression and anxiety symptoms in adolescents and young adults during the COVID-19 pandemic. *Psychol Med.* 2022;52:3222-3230.
10. Belfer ML. Child and adolescent mental disorders: the magnitude of the problem across the globe. *J Child Psychol Psychiatry.* 2008;49:226-36.
11. Rocha TB, Graeff-Martins AS, Kieling C, Rohde LA. Provision of mental healthcare for children and adolescents: a worldwide view. *Curr Opin Psychiatry.* 2015;28:330-5.
12. Park-Lee E, Lipari RN, Hedden SL, Copello E, Kroutil L. Receipt of Services for Substance Use and Mental Health Issues Among Adults: Results from the 2015 National Survey on Drug Use and Health. Rockville: NSDUH; Data Review.

13. Collins KA, Westra HA, Dozois DJ, Burns DD. Gaps in accessing treatment for anxiety and depression: challenges for the delivery of care. *Clin Psychol Rev.* 2004;24:583-616.
14. Mol M, van Genugten C, Dozeman E, van Schaik DJF, Draisma S, Riper H, et al. Why Uptake of Blended Internet-Based Interventions for Depression Is Challenging: A Qualitative Study on Therapists' Perspectives. *J Clin Med.* 2019;9:91.
15. Marcu G, Ondersma SJ, Spiller AN, Broderick BM, Kadri R, Buis LR. The Perceived Benefits of Digital Interventions for Behavioral Health: Qualitative Interview Study. *J Med Internet Res.* 2022;24(3):e34300.
16. Andrews G, Basu A, Cuijpers P, Craske MG, McEvoy P, English CL, et al. Computer therapy for the anxiety and depression disorders is effective, acceptable and practical healthcare: An updated meta-analysis. *J Anxiety Disord.* 2018;55:70-78.
17. Karyotaki E, Riper H, Twisk J, Hoogendoorn A, Kleiboer A, Mira A, et al. Efficacy of Self-guided Internet-Based Cognitive Behavioral Therapy in the Treatment of Depressive Symptoms: A Meta-analysis of Individual Participant Data. *JAMA Psychiatry.* 2017;74:351-359.
18. National Institute for Health and Care Excellence. (2023). Guided self-help digital cognitive behavioural therapy for children and young people with mild to moderate symptoms of anxiety or low mood: early value assessment. <https://www.nice.org.uk/guidance/hte3>
19. March S, Spence SH, Donovan CL, Kenardy JA. Large-Scale Dissemination of Internet-Based Cognitive Behavioral Therapy for Youth Anxiety: Feasibility and Acceptability Study. *J Med Internet Res.* 2018;20(7):e234.
20. Ebert DD, Zarski AC, Christensen H, Stikkelbroek Y, Cuijpers P, Berking M, et al. Internet and computer-based cognitive behavioral therapy for anxiety and depression in youth: a meta-analysis of randomized controlled outcome trials. *PLoS One.* 2015;10(3):e0119895.
21. Sethi S. Treating Youth Depression and Anxiety: A Randomised Controlled Trial Examining the Efficacy of Computerised versus Face-to-face Cognitive Behaviour Therapy. *Aust Psychol.* 2013;48(4):249-257.

22. Vigerland S, Ljótsson B, Gustafsson FB, Hagert S, Thulin U, Andersson G, et al. Attitudes towards the use of computerized cognitive behavior therapy (cCBT) with children and adolescents: A survey among Swedish mental health professionals. *Internet Interv.* 2014;1(3):111-117.
23. MacDonell KW, Prinz RJ. A Review of Technology-Based Youth and Family-Focused Interventions. *Clin Child Fam Psychol Rev.* 2017;20(2):185-200.
24. Wentzel J, van der Vaart R, Bohlmeijer ET, van Gemert-Pijnen JEW. Mixing Online and Face-to-Face Therapy: How to Benefit From Blended Care in Mental Health Care. *JMIR Ment Health.* 2016;3(1):e9.
25. Kleiboer A, Smit J, Bosmans J, Ruwaard J, Andersson G, Topooco N, et al. European COMPARative Effectiveness research on blended Depression treatment versus treatment-as-usual (E-COMPARED): study protocol for a randomized controlled, non-inferiority trial in eight European countries. *Trials.* 2016;17(1):387.
26. Mathiasen K, Andersen TE, Riper H, Kleiboer AA, Roessler KK. Blended CBT versus face-to-face CBT: a randomised non-inferiority trial. *BMC Psychiatry.* 2016;16(1):432.
27. Sekhon M, Cartwright M, Francis JJ. Acceptability of healthcare interventions: an overview of reviews and development of a theoretical framework. *BMC Health Serv Res.* 2017;17(1):88.
28. Becker EM, Jensen-Doss A. Computer-assisted therapies: examination of therapist-level barriers to their use. *Behav Ther.* 2013;44(4):614-24.
29. Schuster R, Pokorny R, Berger T, Topooco N, Laireiter AR. The Advantages and Disadvantages of Online and Blended Therapy: Survey Study Amongst Licensed Psychotherapists in Austria. *J Med Internet Res.* 2018;20(12):e11007.
30. Mendes-Santos C, Weiderpass E, Santana R, Andersson G. Portuguese Psychologists' Attitudes Toward Internet Interventions: Exploratory Cross-Sectional Study. *JMIR Ment Health.* 2020;7(4):e16817.
31. Bruno R, Abbott JM. Australian health professionals' attitudes toward and frequency of use of internet supported psychological interventions. *Int J Ment Health.* 2015 Jun;44(1-2):107–123.

32. Topooco N, Riper H, Araya R, Berking M, Brunn M, Chevreur K, et al. Attitudes towards digital treatment for depression: A European stakeholder survey. *Internet Interv.* 2017;8:1-9.
33. Merkel S. Umsetzungsbarrieren bei der Akzeptanz, Implementation und Verbreitung von Telecare und Telehealth - Ergebnisse einer internationalen Literaturstudie. In: *E-Health-Ökonomie*. Springer Fachmedien Wiesbaden; 2017:117-123.
34. Titzler I, Saruhanjan K, Berking M, Riper H, Ebert DD. Barriers and facilitators for the implementation of blended psychotherapy for depression: A qualitative pilot study of therapists' perspective. *Internet Interv.* 2018;12:150-164.
35. van der Vaart R, Witting M, Riper H, Kooistra L, Bohlmeijer ET, van Gemert-Pijnen LJ. Blending online therapy into regular face-to-face therapy for depression: content, ratio and preconditions according to patients and therapists using a Delphi study. *BMC Psychiatry.* 2014;14:355.
36. Cohen L, Sargent M, Sechrest L. Use of psychotherapy research by professional psychologists. *Am Psychol.* 1986;41(2):198-206.
37. Safran JD, Abrue I, Ogilvie J, DeMaria A. Does psychotherapy influence the clinical practice of researcher-clinicians? *Clin Psychol Sci Pract.* 2011;18:357-371.
38. Alkhalidi B, Sahama T, Huxley C, Gajanayake R. Barriers to implementing eHealth: a multi-dimensional perspective. *Stud Health Technol Inform.* 2014;205:875-9.
39. Lilienfeld SO, Ritschel LA, Lynn SJ, Cautin RL, Lutzman RD. Why many clinical psychologists are resistant to evidence-based practice: Root causes and constructive remedies. *Clin Psychol Rev.* 2013;33(7):883-900.
40. Venkatesh V, Morris MG, Davis GB, Davis FD. User acceptance of information technology: Toward a unified view. *MIS Quarterly.* 2003;27(3):425-478.
41. Al-Mamary YH, Al-nashmi M, Abdul Y, Hassan G, Shamsuddin A. A critical review of models and theories in the field of individual acceptance of technology. *Int J Hybrid Inf Technol.* 2016;9:143-158.

42. Aarons GA. Mental health provider attitudes toward adoption of evidence-based practice: The Evidence-Based Practice Attitude Scale (EBPAS). *Ment Health Serv Res.* 2004;6(2):61-74.
43. Tennant A, Conaghan PG. The Rasch measurement model in rheumatology: What is it and why use it? When should it be applied, and what should one look for in a Rasch paper? *Arthritis Care Res.* 2007;57(8):1358-62.
44. Sasangohar F, Bradshaw MR, Carlson MM, Flack JN, Fowler JC, Freeland D, et al. Adapting an outpatient psychiatric clinic to telehealth during the COVID-19 pandemic: A practice perspective. *J Med Internet Res.* 2020;22(10):e22523.
45. Ohannessian R, Duong TA, Odone A. Global telemedicine implementation and integration within health systems to fight the COVID-19 pandemic: A call to action. *JMIR Public Health Surveill.* 2020;6(2):e18810.
46. Baumeister H, Terhorst Y, Grässle C, Freudenstein M, Nübling R, Ebert DD. Impact of an acceptance-facilitating intervention on psychotherapists' acceptance of blended therapy. *PLoS One.* 2020;15(8):e0236995.
47. Phillips EA, Himmler S, Schreyögg J. Preferences of psychotherapists for blended care in Germany: A discrete choice experiment. *BMC Psychiatry.* 2022;22(1):112.
48. Wind TR, Rijkeboer M, Andersson G, Riper H. The COVID-19 pandemic: The 'black swan' for mental health care and a turning point for e-health. *Internet Interv.* 2020;20:100317.
49. Fishman GD, Elkins J. COVID-19 lessons from the field: Toward a pediatric physical therapy telehealth framework. *Int J Telerehabil.* 2022;14(1):e6448.
50. Ray KN, Ashcraft LE, Mehrotra A, Miller E, Kahn JM. Family perspectives on telemedicine for pediatric subspecialty care. *Telemed J E Health.* 2017;23(10):852-862.
51. Philippi P, Baumeister H, Apolinário-Hagen J, Ebert DD, Hennemann S, Kott L, et al. Acceptance towards digital health interventions - Model validation and further development of the Unified Theory of Acceptance and Use of Technology. *Internet Interv.* 2021;26:100459.
52. Pelckmans M, Nijmolen P, Bloemen M, Kuijpers E, Meershhoek A, Rameckers E, et al. Barriers and facilitators to the acceptance of eHealth interventions

by Dutch pediatric physical therapists in times of the COVID-19 pandemic: A mixed-methods approach. *Pediatr Phys Ther.* 2023;35(2):243-250.

53. De Veirman AEM, Thewissen V, Spruijt MG, Bolman CAW. Factors associated with intention and use of e-mental health by mental health counselors in general practices: Web-based survey. *JMIR Form Res.* 2022;6(12):e34754.

54. van der Vaart R, Atema V, Evers AW. Guided online self-management interventions in primary care: A survey on use, facilitators, and barriers. *BMC Fam Pract.* 2016;17:27.

55. Hoerger M. Participant dropout as a function of survey length in internet-mediated university studies: implications for study design and voluntary participation in psychological research. *Cyberpsychol Behav Soc Netw.* 2010;13(6):697-700.